



[Go to Product page](#)

Datasheet for ABIN7169190

anti-TAO Kinase 1 (TAOK1) (AA 400-659) antibody (HRP)

Overview

| | |
|----------------------|----------------------|
| Quantity: | 100 µL |
| Target: | TAO Kinase 1 (TAOK1) |
| Binding Specificity: | AA 400-659 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | HRP |
| Application: | ELISA |

Product Details

| | |
|-------------------|--|
| Immunogen: | Recombinant Human Serine/threonine-protein kinase TAO1 protein (400-659AA) |
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Purification: | >95%, Protein G purified |

Target Details

| | |
|-------------------|---|
| Target: | TAO Kinase 1 (TAOK1) |
| Alternative Name: | TAOK1 (TAOK1 Products) |
| Background: | Background: Serine/threonine-protein kinase involved in various processes such as p38/MAPK14 stress-activated MAPK cascade, DNA damage response and regulation of |

Target Details

cytoskeleton stability. Phosphorylates MAP2K3, MAP2K6 and MARK2. Acts as an activator of the p38/MAPK14 stress-activated MAPK cascade by mediating phosphorylation and subsequent activation of the upstream MAP2K3 and MAP2K6 kinases. Involved in G-protein coupled receptor signaling to p38/MAPK14. In response to DNA damage, involved in the G2/M transition DNA damage checkpoint by activating the p38/MAPK14 stress-activated MAPK cascade, probably by mediating phosphorylation of MAP2K3 and MAP2K6. Acts as a regulator of cytoskeleton stability by phosphorylating 'Thr-208' of MARK2, leading to activate MARK2 kinase activity and subsequent phosphorylation and detachment of MAPT/TAU from microtubules. Also acts as a regulator of apoptosis: regulates apoptotic morphological changes, including cell contraction, membrane blebbing and apoptotic bodies formation via activation of the MAPK8/JNK cascade.

Aliases: 2810468K05Rik antibody, AU020252 antibody, D130018F14Rik antibody, EC 2.7.11.1 antibody, FLJ14314 antibody, hKFC B antibody, hKFC-B antibody, KIAA1361 antibody, Kinase from chicken homolog B antibody, MAP3K16 antibody, MARK kinase antibody, MARKK antibody, MGC29021 antibody, Microtubule affinity regulating kinase kinase antibody, mKIAA1361 antibody, Prostate-derived sterile 20-like kinase 2 antibody, PSK2 antibody, Serine/threonine kinase TAO1 antibody, Serine/threonine protein kinase TAO1 antibody, Serine/threonine protein kinase TAO1 homolog antibody, Serine/threonine-protein kinase TAO1 antibody, STE20 like kinase antibody, STE20 like kinase PSK2 antibody, STE20-like kinase PSK2 antibody, TAO kinase 1 antibody, Taok1 antibody, TAOK1_HUMAN antibody, Thousand and one amino acid protein 1 antibody

UniProt: [Q7L7X3](#)

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be

Handling

handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.