

Datasheet for ABIN7169368
anti-SAA2 antibody (AA 19-94)[Go to Product page](#)

2 Images

Overview

Quantity:	100 µg
Target:	SAA2
Binding Specificity:	AA 19-94
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SAA2 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), ELISA

Product Details

Immunogen:	Recombinant Human Serum amyloid A-2 protein (19-94AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	SAA2
Alternative Name:	SAA2 (SAA2 Products)
Background:	Background: Major acute phase reactant. Apolipoprotein of the HDL complex. Aliases: SAA2 antibody, Serum amyloid A-2 protein antibody, SAA2 antibody

Target Details

UniProt:	P0DJ19
Pathways:	Toll-Like Receptors Cascades


Application Details

Application Notes:	Recommended dilution: IHC:1:500-1:1000,
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Immunohistochemistry

Image 1. IHC image of ABIN7169368 diluted at 1:600 and staining in paraffin-embedded human liver cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunohistochemistry

Image 2. IHC image of ABIN7169368 diluted at 1:600 and staining in paraffin-embedded human liver tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.