antibodies -online.com





Datasheet for ABIN716942 anti-ACPT antibody (AA 65-160) (Cy7)

Go to Product page

| () | 11/0 | K\ / | iew | 1 |
|-----|--------|--------|---------------------|---|
| | \cup | 'I V/I | $\square \vee \vee$ | ı |
| | | | | |

| Quantity: | 100 μL |
|----------------------|---|
| Target: | ACPT |
| Binding Specificity: | AA 65-160 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This ACPT antibody is conjugated to Cy7 |
| Application: | Flow Cytometry (FACS), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Western Blotting (WB) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human ACPT |
|-----------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Predicted Reactivity: | Mouse,Rat,Dog,Horse |
| Purification: | Purified by Protein A. |

Target Details

| Target: | ACPT |
|-------------------|----------------------|
| Alternative Name: | ACPT (ACPT Products) |

Target Details

| Background: | Synonyms: Acid phosphatase testicular, Testicular acid phosphatase, PPAT_HUMAN. | | |
|---------------------|---|--|--|
| | Background: Acid phosphatases are enzymes capable of hydrolyzing orthophosphoric acid | | |
| | esters in an acid medium. This gene is upregulated by androgens and is down-regulated by | | |
| | estrogens in the prostate cancer cell line. ACPT dephosphorylates receptor tyrosine-protein kinase erbB-4 and inhibits the ligand-induced proteolytic cleavage. ACPT is predominantly | | |
| | | | |
| | expressed in testis but exhibits a lower level of expression in testicular cancer tissues than in | | |
| | normal tissues. ACPT has structural similarity to prostatic and lysosomal acid phosphatases. | | |
| | Alternatively spliced transcript variants have been described, but their biological validity has not | | |
| | been determined. | | |
| Gene ID: | 93650 | | |
| Application Details | | | |
| Application Notes: | FCM 1:20-100 | | |
| | IF(IHC-P) 1:50-200 | | |
| | IF(IHC-F) 1:50-200 | | |
| | IF(ICC) 1:50-200 | | |
| Restrictions: | For Research Use only | | |
| Handling | | | |
| Format: | Liquid | | |
| Concentration: | 1 μg/μL | | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. | | |
| Preservative: | ProClin | | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. | | |
| Storage: | -20 °C | | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | | |
| Expiry Date: | 12 months | | |
| | | | |