antibodies -online.com





anti-SH2D1B antibody (AA 1-77) (Biotin)



Overview

Quantity:	100 μg
Target:	SH2D1B
Binding Specificity:	AA 1-77
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SH2D1B antibody is conjugated to Biotin
Application:	ELISA

Product Details

Immunogen:	Recombinant Human SH2 domain-containing protein 1B protein (1-77AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	SH2D1B
Alternative Name:	SH2D1B (SH2D1B Products)
Background:	Background: Cytoplasmic adapter regulating receptors of the signaling lymphocytic activation
	molecule (SLAM) family such as CD84, SLAMF1, LY9 and CD244 (PubMed:11689425). In SLAM

signaling seems to cooperate with SH2D1A/SAP. Plays a role in regulation of effector functions of natural killer (NK) cells by controlling signal transduction through CD244/2B4 without effecting its tyrosine phosphorylation, downstream signaling involves PLCG1 and ERK activation (PubMed:24687958). Activation of SLAMF7-mediated NK cell function does not effect receptor tyrosine phosphorylation but distal signaling (By similarity). In the context of NK cell-mediated cytotoxicity does not enhance conjugate formation with target cells but stimulates polarization of the microtubule-organizing center and cytotoxic granules toward the NK cell synapse (PubMed:24687958). Negatively regulates CD40-induced cytokine procuction dendritic cells downstream of SLAM family receptors probably by inducing activation of the PI3K pathway to inhibit p38 MAPK and JNK activation (By similarity). Aliases: SH2D1B antibody, EAT2 antibody, SH2 domain-containing protein 1B antibody,

EWS/FLI1-activated transcript 2 antibody, EAT-2 antibody

UniProt: 014796

Pathways: Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process

Application Details

Optimal working dilution should be determined by the investigator. Application Notes:

For Research Use only

Handling

Restrictions:

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.