

Datasheet for ABIN7169677
anti-STAT3 antibody (AA 50-240) (FITC)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	STAT3
Binding Specificity:	AA 50-240
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This STAT3 antibody is conjugated to FITC
Application:	Please inquire

Product Details

Immunogen:	Recombinant Human Signal transducer and activator of transcription 3 protein (50-240AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	STAT3
Alternative Name:	STAT3 (STAT3 Products)
Background:	Background: Signal transducer and transcription activator that mediates cellular responses to interleukins, KITLG/SCF, LEP and other growth factors. Once activated, recruits coactivators,

Target Details

such as NCOA1 or MED1, to the promoter region of the target gene (PubMed:17344214). May mediate cellular responses to activated FGFR1, FGFR2, FGFR3 and FGFR4. Binds to the interleukin-6 (IL-6)-responsive elements identified in the promoters of various acute-phase protein genes. Activated by IL31 through IL31RA. Involved in cell cycle regulation by inducing the expression of key genes for the progression from G1 to S phase, such as CCND1 (PubMed:17344214). Mediates the effects of LEP on melanocortin production, body energy homeostasis and lactation (By similarity). May play an apoptotic role by transactivating BIRC5 expression under LEP activation (PubMed:18242580). Cytoplasmic STAT3 represses macroautophagy by inhibiting EIF2AK2/PKR activity.

Aliases: 1110034C02Rik antibody, Acute Phase Response Factor antibody, Acute-phase response factor antibody, ADMIO antibody, APRF antibody, AW109958 antibody, DNA binding protein APRF antibody, FLJ20882 antibody, HIES antibody, MGC16063 antibody, Signal transducer and activator of transcription 3 (acute phase response factor) antibody, Signal transducer and activator of transcription 3 antibody, STAT 3 antibody, Stat3 antibody, STAT3_HUMAN antibody

UniProt:	P40763
Pathways:	JAK-STAT Signaling , RTK Signaling , Interferon-gamma Pathway , Neurotrophin Signaling Pathway , Dopaminergic Neurogenesis , Response to Growth Hormone Stimulus , Carbohydrate Homeostasis , Stem Cell Maintenance , Hepatitis C , Protein targeting to Nucleus , Feeding Behaviour , CXCR4-mediated Signaling Events , Signaling of Hepatocyte Growth Factor Receptor

Application Details

Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C

Handling

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.