antibodies -online.com





anti-SLC38A9 antibody (AA 1-119)





Overview

Overview	
Quantity:	100 μg
Target:	SLC38A9
Binding Specificity:	AA 1-119
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SLC38A9 antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF)
Product Details	
Immunogen:	Recombinant Human Sodium-coupled neutral amino acid transporter 9 protein (1-119AA)
Isotype:	IgG

Target Details

Cross-Reactivity:

Purification:

Human

>95%, Protein G purified

Target:	SLC38A9
Alternative Name:	SLC38A9 (SLC38A9 Products)
Background:	Background: Lysosomal amino acid transporter involved in the activation of mTORC1 in
	response to amino acids. Probably acts as an amino acid sensor of the Rag GTPases and

Ragulator complexes, 2 complexes involved in amino acid sensing and activation of mTORC1, a signaling complex promoting cell growth in response to growth factors, energy levels, and amino acids (PubMed:25561175, PubMed:25567906). Following activation by amino acids, the Ragulator and Rag GTPases function as a scaffold recruiting mTORC1 to lysosomes where it is in turn activated. SLC38A9 mediates transport of amino acids with low capacity and specificity with a slight preference for polar amino acids, suggesting that it acts as an amino acid sensor instead (PubMed:25561175, PubMed:25567906). The high concentration of arginine in lysosomes suggests that it acts as an arginine sensor (PubMed:25567906).


Aliases: FLJ46104 antibody, FLJ90709 antibody, Homo sapiens solute carrier family 38, member 9 antibody, MGC120544 antibody, OTTHUMP00000161835 antibody, Putative sodium coupled neutral amino acid transporter 9 antibody, Putative sodium-coupled neutral amino acid transporter 9 antibody, Putative uncharacterized protein FLJ90709 antibody, S38A9_HUMAN antibody, SLC38A9 antibody

UniProt:

Q8NBW4


Application Details

Application Notes:	Recommended dilution: WB:1:500-1:5000, IHC:1:20-1:200, IF:1:50-1:200,
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.




Immunofluorescence

Image 1. Immunofluorescent analysis of U251 cells using ABIN7169972 at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)



Western Blotting

Image 2. Western Blot Positive WB detected in: Jurkat whole cell lysate All lanes: SLC38A9 antibody at 3.5 μg/mL Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 64, 53, 57 kDa Observed band size: 64 kDa



Immunohistochemistry

Image 3. Immunohistochemistry of paraffin-embedded human liver tissue using ABIN7169972 at dilution of 1:100

Please check the product details page for more images. Overall 4 images are available for ABIN7169972.