

[Go to Product page](#)

Datasheet for ABIN7170133  
**anti-SLC22A1 antibody (AA 43-149) (FITC)**

## Overview

Quantity:	100 µg
Target:	SLC22A1
Binding Specificity:	AA 43-149
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SLC22A1 antibody is conjugated to FITC
Application:	Please inquire

## Product Details

Immunogen:	Recombinant Human Solute carrier family 22 member 1 protein (43-149AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

## Target Details

Target:	SLC22A1
Alternative Name:	SLC22A1 ( <a href="#">SLC22A1 Products</a> )
Background:	Background: Translocates a broad array of organic cations with various structures and molecular weights including the model compounds 1-methyl-4-phenylpyridinium (MPP),

## Target Details

tetraethylammonium (TEA), N-1-methylnicotinamide (NMN), 4-(4-(dimethylamino)styryl)-N-methylpyridinium (ASP), the endogenous compounds choline, guanidine, histamine, epinephrine, adrenaline, noradrenaline and dopamine, and the drugs quinine, and metformin. The transport of organic cations is inhibited by a broad array of compounds like tetramethylammonium (TMA), cocaine, lidocaine, NMDA receptor antagonists, atropine, prazosin, cimetidine, TEA and NMN, guanidine, cimetidine, choline, procainamide, quinine, tetrabutylammonium, and tetrapentylammonium. Translocates organic cations in an electrogenic and pH-independent manner. Translocates organic cations across the plasma membrane in both directions. Transports the polyamines spermine and spermidine. Transports pramipexole across the basolateral membrane of the proximal tubular epithelial cells. The choline transport is activated by MMTS. Regulated by various intracellular signaling pathways including inhibition by protein kinase A activation, and endogenous activation by the calmodulin complex, the calmodulin-dependent kinase II and LCK tyrosine kinase.

Aliases: hOCT1 antibody, OCT1 antibody, oct1\_cds antibody, Organic cation transporter 1 antibody, S22A1\_HUMAN antibody, Slc22a1 antibody, solute carrier family 22 (organic cation transporter), member 1 antibody, Solute carrier family 22 member 1 antibody

UniProt: [O15245](#)

Pathways: [Hormone Transport](#)

## Application Details

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C, -80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.