

[Go to Product page](#)

Datasheet for ABIN7170401  
**anti-SNX5 antibody (AA 7-245) (HRP)**

## Overview

Quantity:	100 µL
Target:	SNX5
Binding Specificity:	AA 7-245
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SNX5 antibody is conjugated to HRP
Application:	ELISA

## Product Details

Immunogen:	Recombinant Human Sorting nexin-5 protein (7-245AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

## Target Details

Target:	SNX5
Alternative Name:	SNX5 ( <a href="#">SNX5 Products</a> )
Background:	Background: Involved in several stages of intracellular trafficking. Interacts with membranes containing phosphatidylinositol 3-phosphate (PtdIns(3P)) or phosphatidylinositol 3,4-

## Target Details

bisphosphate (PtdIns(3,4)P<sub>2</sub>) (PubMed:15561769). Acts in part as component of the retromer membrane-deforming SNX-BAR subcomplex. The SNX-BAR retromer mediates retrograde transport of cargo proteins from endosomes to the trans-Golgi network (TGN) and is involved in endosome-to-plasma membrane transport for cargo protein recycling. The SNX-BAR subcomplex functions to deform the donor membrane into a tubular profile called endosome-to-TGN transport carrier (ETC) (Probable). Does not have in vitro vesicle-to-membrane remodeling activity (PubMed:23085988). Involved in retrograde transport of lysosomal enzyme receptor IGF2R (PubMed:17148574, PubMed:18596235). May function as link between endosomal transport vesicles and dynactin (Probable). Plays a role in the internalization of EGFR after EGF stimulation (Probable). Involved in EGFR endosomal sorting and degradation, the function involves PIP5K1C isoform 3 and is retromer-independent (PubMed:23602387). Together with PIP5K1C isoform 3 facilitates HGS interaction with ubiquitinated EGFR, which initiates EGFR sorting to intraluminal vesicles (ILVs) of the multivesicular body for subsequent lysosomal degradation (Probable). Involved in E-cadherin sorting and degradation, inhibits PIP5K1C isoform 3-mediated E-cadherin degradation (PubMed:24610942). Plays a role in macropinocytosis (PubMed:18854019, PubMed:21048941).

Aliases: 0910001N05Rik antibody, 1810032P22Rik antibody, AU019504 antibody, D2Ert52e antibody, FLJ10931 antibody, OTTHUMP00000030340 antibody, OTTHUMP00000062927 antibody, OTTMUSP00000003367 antibody, OTTMUSP00000003368 antibody, RP11-504H3.2 antibody, RP23-35E16.2 antibody, snoRNA MBI-43 antibody, SNX 5 antibody, SNX5 antibody, SNX5\_HUMAN antibody, Sorting nexin 5 antibody, Sorting nexin-5 antibody

UniProt: [Q9Y5X3](#)

## Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be

## Handling

---

handled by trained staff only.

---

Storage: -20 °C,-80 °C

---

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.