

Datasheet for ABIN7170456
anti-SPEF2 antibody (AA 932-1230)[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	SPEF2
Binding Specificity:	AA 932-1230
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SPEF2 antibody is un-conjugated
Application:	ELISA, Immunofluorescence (IF), Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human Sperm flagellar protein 2 protein (932-1230aa)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	SPEF2
Alternative Name:	SPEF2 (SPEF2 Products)
Background:	Background: Required for correct axoneme development. Aliases: Sperm flagellar protein 2 (Protein KPL2), SPEF2, KIAA1770 KPL2

Target Details

UniProt: [Q9C093](#)

Application Details

Application Notes: Recommended dilution: IHC:1:100-1:500, IF:1:100-1:500,

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4


Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C


Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Immunohistochemistry

Image 1. IHC image of ABIN7170456 diluted at 1:200 and staining in paraffin-embedded human kidney tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.05 % DAB.



Immunofluorescence

Image 2. Immunofluorescence staining of MCF-7 cells with ABIN7170456 at 1:100, counter-stained with DAPI. The cells were fixed in 4 % formaldehyde and blocked in 10 % normal Goat Serum. The cells were then incubated with the antibody overnight at 4 °C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).