
antibodies - online.com







anti-SDHA antibody (AA 44-293)





Overview	
Quantity:	100 μg
Target:	SDHA
Binding Specificity:	AA 44-293
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SDHA antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF)
Product Details	
Immunogon:	Pacambinant Human Susainata dabudraganasa (ubiguinana) flavanratain subunit

Immunogen:	Recombinant Human Succinate dehydrogenase [ubiquinone] flavoprotein subunit, mitochondrial protein (44-293AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	SDHA
Alternative Name:	SDHA (SDHA Products)
Background:	Background: Flavoprotein (FP) subunit of succinate dehydrogenase (SDH) that is involved in

complex II of the mitochondrial electron transport chain and is responsible for transferring electrons from succinate to ubiquinone (coenzyme Q).

Aliases: CMD1GG antibody, DHSA_HUMAN antibody, Flavoprotein subunit of complex II antibody, Fp antibody, PGL5 antibody, SDH 2 antibody, SDH1 antibody, SDH2 antibody, SDHA antibody, SDHF antibody, Succinate dehydrogenase [ubiquinone] flavoprotein subunit antibody, Succinate dehydrogenase [ubiquinone] flavoprotein subunit mitochondrial antibody, Succinate dehydrogenase [ubiquinone] flavoprotein subunit, mitochondrial antibody, Succinate dehydrogenase complex flavoprotein subunit A antibody, Succinate dehydrogenase complex flavoprotein subunit precursor antibody, Succinate dehydrogenase complex subunit A antibody, Succinate dehydrogenase complex subunit A flavoprotein (Fp) antibody, Succinate dehydrogenase complex subunit A flavoprotein antibody

UniProt:

P31040

Application Details

Application Notes:	Recommended dilution: WB:1:500-1:5000, IHC:1:20-1:200, IF:1:50-1:200,
Restrictions:	For Research Use only
Handling	
Format:	Liquid

Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.




Immunocytochemistry

Image 1. Immunocytochemistry analysis of human skeletal muscle tissue using ABIN7170882 at dilution of 1:100



Western Blotting

Image 2. Western blot All lanes: SDHA antibody at $2 \mu g/mL + 293T$ whole cell lysate Secondary Goat polyclonal to rabbit lgG at 1/15000 dilution Predicted band size: 73, 68, 57 kDa Observed band size: 73, 57 kDa



Immunofluorescence

Image 3. Immunofluorescent analysis of PC-3 cells using ABIN7170882 at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)

Please check the product details page for more images. Overall 4 images are available for ABIN7170882.