antibodies .- online.com





anti-SUN2 antibody (AA 45-155) (Biotin)



Go to Product page

()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μg
Target:	SUN2
Binding Specificity:	AA 45-155
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SUN2 antibody is conjugated to Biotin
Application:	ELISA

Product Details

Immunogen:	Recombinant Human SUN domain-containing protein 2 protein (45-155AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	SUN2
Alternative Name:	SUN2 (SUN2 Products)
Background:	Background: As a component of the LINC (Linker of Nucleoskeleton and Cytoskeleton)
	complex, involved in the connection between the nuclear lamina and the cytoskeleton. The

nucleocytoplasmic interactions established by the LINC complex play an important role in the transmission of mechanical forces across the nuclear envelope and in nuclear movement and positioning. Specifically, SYNE2 and SUN2 assemble in arrays of transmembrane actinassociated nuclear (TAN) lines which are bound to F-actin cables and couple the nucleus to retrograde actin flow during actin-dependent nuclear movement. Required for interkinetic nuclear migration (INM) and essential for nucleokinesis and centrosome-nucleus coupling during radial neuronal migration in the cerebral cortex and during glial migration. Required for nuclear migration in retinal photoreceptor progenitors implicating association with cytoplasmic dynein-dynactin and kinesin motor complexes, and probably B-type lamins, SUN1 and SUN2 seem to act redundantly. The SUN1/2:KASH5 LINC complex couples telomeres to microtubules during meiosis, SUN1 and SUN2 seem to act at least partial redundantly. Anchors chromosome movement in the prophase of meiosis and is involved in selective gene expression of coding and non-coding RNAs needed for gametogenesis. Required for telomere attachment to nuclear envelope and gametogenesis. May also function on endocytic vesicles as a receptor for RAB5-GDP and participate in the activation of RAB5.

Aliases: SUN2 antibody, FRIGG antibody, KIAA0668 antibody, RAB5IP antibody, UNC84B antibody, SUN domain-containing protein 2 antibody, Protein unc-84 homolog B antibody, Rab5-interacting protein antibody, Rab5IP antibody, Sad1/unc-84 protein-like 2 antibody

UniProt:

Q9UH99

Pathways:

Maintenance of Protein Location, SARS-CoV-2 Protein Interactome, The Global Phosphorylation Landscape of SARS-CoV-2 Infection

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.	
Restrictions:	For Research Use only	

Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling

Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.