antibodies .- online.com





anti-SMARCC2 antibody (AA 300-647) (FITC)



Go to Product page

\sim			
	N/P	r\/	i⊢₩

Quantity:	100 μL
Target:	SMARCC2
Binding Specificity:	AA 300-647
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SMARCC2 antibody is conjugated to FITC
Application:	Please inquire

Product Details

Immunogen:	Recombinant Human SWI/SNF complex subunit SMARCC2 protein (300-647aa)	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	>95%, Protein G purified	

Target Details

Target:	SMARCC2	
Alternative Name:	SMARCC2 (SMARCC2 Products)	
Background:	Background: Background: Involved in transcriptional activation and repression of select genes by chrom remodeling (alteration of DNA-nucleosome topology). Can stimulate the ATPase activity of	

catalytic subunit of these complexes. May be required for CoREST dependent repression of neuronal specific gene promoters in non-neuronal cells. Belongs to the neural progenitors-specific chromatin remodeling complex (npBAF complex) and the neuron-specific chromatin remodeling complex (nBAF complex). During neural development a switch from a stem/progenitor to a post-mitotic chromatin remodeling mechanism occurs as neurons exit the cell cycle and become committed to their adult state. The transition from proliferating neural stem/progenitor cells to post-mitotic neurons requires a switch in subunit composition of the npBAF and nBAF complexes. As neural progenitors exit mitosis and differentiate into neurons, npBAF complexes which contain ACTL6A/BAF53A and PHF10/BAF45A, are exchanged for homologous alternative ACTL6B/BAF53B and DPF1/BAF45B or DPF3/BAF45C subunits in neuron-specific complexes (nBAF). The npBAF complex is essential for the self-renewal/proliferative capacity of the multipotent neural stem cells. The nBAF complex along with CREST plays a role regulating the activity of genes essential for dendrite growth (By similarity).

Aliases: SWI/SNF complex subunit SMARCC2 (BRG1-associated factor 170) (BAF170) (SWI/SNF complex 170 kDa subunit) (SWI/SNF-related matrix-associated actin-dependent regulator of chromatin subfamily C member 2), SMARCC2, BAF170

UniProt:

Q8TAQ2

Application Details

Restrictions:

For Research Use only

Handling

Format:	Liquid	
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C,-80 °C	
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.	