
antibodies -online.com





anti-SMARCD1 antibody (AA 7-124)





Go to Product page

\sim					
()	VE	۲۱	/1	\triangle	Λ

Overview		
Quantity:	100 μg	
Target:	SMARCD1	
Binding Specificity:	AA 7-124	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This SMARCD1 antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF)	
Product Details		
Immunogen:	Recombinant Human SWI/SNF-related matrix-associated actin-dependent regulator of	
	chromatin subfamily D member 1 protein (7-124AA)	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	>95%, Protein G purified	
Target Details		
Target:	SMARCD1	
Alternative Name:	SMARCD1 (SMARCD1 Products)	
Background:	Background: Involved in chromatin remodeling. Belongs to the neural progenitors-specific	

chromatin remodeling complex (npBAF complex) and the neuron-specific chromatin remodeling complex (nBAF complex). During neural development a switch from a stem/progenitor to a post-mitotic chromatin remodeling mechanism occurs as neurons exit the cell cycle and become committed to their adult state. The transition from proliferating neural stem/progenitor cells to post-mitotic neurons requires a switch in subunit composition of the npBAF and nBAF complexes. As neural progenitors exit mitosis and differentiate into neurons, npBAF complexes which contain ACTL6A/BAF53A and PHF10/BAF45A, are exchanged for homologous alternative ACTL6B/BAF53B and DPF1/BAF45B or DPF3/BAF45C subunits in neuron-specific complexes (nBAF). The npBAF complex is essential for the selfrenewal/proliferative capacity of the multipotent neural stem cells. The nBAF complex along with CREST plays a role regulating the activity of genes essential for dendrite growth (By similarity). Has a strong influence on vitamin D-mediated transcriptional activity from an enhancer vitamin D receptor element (VDRE). May be a link between mammalian SWI-SNF-like chromatin remodeling complexes and the vitamin D receptor (VDR) heterodimer. Mediates critical interactions between nuclear receptors and the BRG1/SMARCA4 chromatin-remodeling complex for transactivation.

Aliases: 60 kDa BRG-1/Brm-associated factor subunit A antibody, 60 kDa BRG1/Brm associated factor subunit A antibody, BAF60A antibody, BRG1 associated factor 60A antibody, BRG1-associated factor 60A antibody, Chromatin remodeling complex BAF60A subunit antibody, CRACD1 antibody, Mammalian chromatin remodeling complex BRG1 associated factor 60A antibody, Rsc6p antibody, SMARCD1 antibody, SMRD1 antibody, SMRD1_HUMAN antibody, SWI/SNF complex 60 kDa subunit A antibody, SWI/SNF complex 60 kDa subunit antibody, SWI/SNF related matrix associated actin dependent regulator of chromatin d1 antibody, SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily d, member 1 antibody, SWI/SNF-related matrix-associated actin-dependent regulator of chromatin subfamily D member 1 antibody, SWp73 like protein antibody

UniProt:

Format:

Q96GM5

Liquid


Application Details

Application Notes:	Recommended dilution: WB:1:500-1:5000, IHC:1:20-1:200, IF:1:50-1:200,	
Restrictions:	For Research Use only	
Handling		

Handling


Buffer:	Preservative: 0.03 % Proclin 300	
	Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C,-80 °C	
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.	

Images




Immunohistochemistry

Image 1. Immunohistochemistry of paraffin-embedded human lymph node tissue using ABIN7171071 at dilution of 1:100



Western Blotting

Image 2. Western Blot Positive WB detected in: Hela whole cell lysate, Jurkat whole cell lysate, HepG2 whole cell lysate, K562 whole cell lysate All lanes: SMARCD1 antibody at 3.2 μ g/mL Secondary Goat polyclonal to rabbit lgG at 1/50000 dilution Predicted band size: 59, 54 kDa Observed band size: 59 kDa



Immunohistochemistry

Image 3. Immunohistochemistry of paraffin-embedded human cervical cancer using ABIN7171071 at dilution of 1:100

Please check the product details page for more images. Overall 4 images are available for ABIN7171071.