antibodies - online.com







anti-CD1e antibody (AA 198-294)



Image



\sim	
()\/△	rview
\cup	1 410 44

Quantity:	100 μg
Target:	CD1e
Binding Specificity:	AA 198-294
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CD1e antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human T-cell surface glycoprotein CD1e, membrane-associated protein (198-294AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	CD1e
Alternative Name:	CD1E (CD1e Products)
Background:	Background: T-cell surface glycoprotein CD1e, soluble binds diacetylated lipids, including

Target Details

phosphatidyl inositides and diacylated sulfoglycolipids, and is required for the presentation of glycolipid antigens on the cell surface. The membrane-associated form is not active.

Aliases: CD_antigen=CD1e antibody, CD1A antibody, CD1e antibody, CD1e antigen antibody, CD1E antigen, e polypeptide antibody, CD1e molecule antibody, CD1E_HUMAN antibody, Differentiation antigen CD1-alpha-3 antibody, hCD1e antibody, Leukocyte differentiation antigen antibody, R2 antibody, R2G1 antibody, sCD1e antibody, soluble antibody, T-cell surface glycoprotein CD1e antibody, T-cell surface glycoprotein CD1e, soluble antibody

UniProt:


P15812

Application Details

Storage Comment:

Application Notes:	Recommended dilution: IHC:1:200-1:500,
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300
	Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be
	handled by trained staff only.
Storage:	-20 °C,-80 °C
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Immunohistochemistry

Image 1. IHC image of ABIN7171377 diluted at 1:300 and staining in paraffin-embedded human endometrial cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.