

Datasheet for ABIN7171574
anti-BTAF1 antibody (AA 1750-1849) (Biotin)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	BTAF1
Binding Specificity:	AA 1750-1849
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BTAF1 antibody is conjugated to Biotin
Application:	ELISA

Product Details

Immunogen:	Recombinant Human TATA-binding protein-associated factor 172 protein (1750-1849AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	BTAF1
Alternative Name:	BTAF1 (BTAF1 Products)
Background:	Background: Regulates transcription in association with TATA binding protein (TBP). Removes TBP from the TATA box in an ATP-dependent manner.

Target Details

Aliases: ATP dependent helicase BTAF1 antibody, ATP-dependent helicase BTAF1 antibody, B-TFIID transcription factor associated 170 kDa subunit antibody, B-TFIID transcription factor associated, 170 kDa (Mot1 homolog, S. cerevisiae) antibody, B-TFIID transcription factor-associated 170 kDa subunit antibody, BTAF 1 RNA polymerase II antibody, BTAF1 antibody, BTAF1 RNA polymerase II B TFIID transcription factor associated 170 kDa antibody, BTAF1_HUMAN antibody, MOT 1 antibody, MOT1 antibody, MOT1 Yeast homolog of antibody, TAF 172 antibody, TAF II 170 antibody, TAF(II)170 antibody, TAF-172 antibody, TAF172 antibody, TAFII170 antibody, TATA binding protein associated factor 172 antibody, TATA-binding protein-associated factor 172 antibody, TBP associated factor 172 antibody, TBP-associated factor 172 antibody

UniProt: [O14981](#)

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.