# antibodies -online.com





# anti-TBC1D23 antibody (AA 510-605)



# Image



#### Overview

Quantity:	100 μg
Target:	TBC1D23
Binding Specificity:	AA 510-605
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TBC1D23 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

### **Product Details**

Immunogen:	Recombinant Human TBC1 domain family member 23 protein (510-605AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

# Target Details

Target:	TBC1D23
Alternative Name:	TBC1D23 (TBC1D23 Products)
Background:	Background: Golgi apparatus
	Aliases: DKFZp667G062 antibody, FLJ11046 antibody, HCV non-structural protein 4A-

#### **Target Details**

	transactivated protein 1 antibody, NS4ATP1 antibody, TBC1 domain family member 23
	antibody, TBC1D23 antibody, TBC23_HUMAN antibody
UniProt:	Q9NUY8


#### **Application Details**

Application Notes:	Recommended dilution: IHC:1:200-1:500,
Restrictions:	For Research Use only

#### Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300
	Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
	σοποιιταστικό, σο το σήσει ση στο τιπτ. Σο, ρ. τ. τ.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be
	handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

#### Images



#### **Immunohistochemistry**

Image 1. IHC image of ABIN7171618 diluted at 1:200 and staining in paraffin-embedded human kidney tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.