antibodies -online.com





Datasheet for ABIN7171734

anti-TRF2 antibody (AA 78-238) (Biotin)



Overview

Quantity:	100 μg
Target:	TRF2 (TERF2)
Binding Specificity:	AA 78-238
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TRF2 antibody is conjugated to Biotin
Application:	ELISA

Product Details

Immunogen:	Recombinant Human Telomeric repeat-binding factor 2 protein (78-238AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	TRF2 (TERF2)
Alternative Name:	TERF2 (TERF2 Products)
Background:	Background: Binds the telomeric double-stranded 5\\\'-TTAGGG-3\\\' repeat and plays a central role in telomere maintenance and protection against end-to-end fusion of chromosomes. In

addition to its telomeric DNA-binding role, required to recruit a number of factors and enzymes required for telomere protection, including the shelterin complex, TERF2IP/RAP1 and DCLRE1B/Apollo. Component of the shelterin complex (telosome) that is involved in the regulation of telomere length and protection. Shelterin associates with arrays of doublestranded 5\\\'-TTAGGG-3\\\' repeats added by telomerase and protects chromosome ends, without its protective activity, telomeres are no longer hidden from the DNA damage surveillance and chromosome ends are inappropriately processed by DNA repair pathways. Together with DCLRE1B/Apollo, plays a key role in telomeric loop (T loop) formation by generating 3\\\' single-stranded overhang at the leading end telomeres: T loops have been proposed to protect chromosome ends from degradation and repair. Required both to recruit DCLRE1B/Apollo to telomeres and activate the exonuclease activity of DCLRE1B/Apollo. Preferentially binds to positive supercoiled DNA. Together with DCLRE1B/Apollo, required to control the amount of DNA topoisomerase (TOP1, TOP2A and TOP2B) needed for telomere replication during fork passage and prevent aberrant telomere topology. Recruits TERF2IP/RAP1 to telomeres, thereby participating in to repressing homology-directed repair (HDR), which can affect telomere length.

Aliases: Telomeric DNA binding protein antibody, Telomeric DNA-binding protein antibody, Telomeric repeat binding factor 2 antibody, Telomeric repeat binding protein 2 antibody, Telomeric repeat-binding factor 2 antibody, TERF 2 antibody, Terf2 antibody, TERF2_HUMAN antibody, TRBF 2 antibody, TRBF2 antibody, TRF2 antibody, TRF2 antibody, TTAGGG repeat binding factor 2 antibody, TTAGGG repeat-binding factor 2 antibody

UniProt:

Q15554

Pathways:

Cell Division Cycle, Telomere Maintenance

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Format:

Buffer:

Preservative: 0.03 % Proclin 300

Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4

Preservative:

ProClin

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.