
antibodies - online.com







anti-TES antibody (AA 130-260)





\sim	
()\/△	rview
\circ	

Quantity:	100 μg
Target:	TES
Binding Specificity:	AA 130-260
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TES antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF)

Product Details

Immunogen:	Recombinant Human Testin protein (130-260AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	TES
Alternative Name:	TES (TES Products)
Background:	Background: Scaffold protein that may play a role in cell adhesion, cell spreading and in the
	reorganization of the actin cytoskeleton. Plays a role in the regulation of cell proliferation. May

Target Details

act as a tumor suppressor.	Inhibits tumor cell growth.

Aliases: DKFZP586B2022 antibody, MGC1146 antibody, TES antibody, TES_HUMAN antibody, TESS 2 antibody, TESS antibody, TESS antibody, Testis derived transcript (3 LIM domains) antibody, Testis derived transcript antibody

UniProt: Q9UGI8


Application Details

Application Notes:	Recommended dilution: WB:1:500-1:5000, IHC:1:500-1:1000, IF:1:200-1:500,
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images




Immunofluorescence

Image 1. Immunofluorescence staining of Hela cells with ABIN7171794 at 1:266, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Western Blotting

Image 2. Western Blot Positive WB detected in: 293 whole cell lysate, Hela whole cell lysate All lanes: TES antibody at 2.92 μg/mL Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 48, 47 kDa Observed band size: 48 kDa



Immunohistochemistry

Image 3. IHC image of ABIN7171794 diluted at 1:800 and staining in paraffin-embedded human liver cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.