# antibodies -online.com





## anti-TRAF3 antibody (AA 1-40) (FITC)



#### Overview

Quantity:	100 μg
Target:	TRAF3
Binding Specificity:	AA 1-40
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TRAF3 antibody is conjugated to FITC
Application:	Please inquire

#### **Product Details**

Immunogen:	Recombinant Human TNF receptor-associated factor 3 protein (1-40AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

### Target Details

Target:	TRAF3
Alternative Name:	TRAF3 (TRAF3 Products)
Background:	Background: Regulates pathways leading to the activation of NF-kappa-B and MAP kinases, and plays a central role in the regulation of B-cell survival. Part of signaling pathways leading to the

production of cytokines and interferon. Required for normal antibody isotype switching from IgM to IgG. Plays a role T-cell dependent immune responses. Plays a role in the regulation of antiviral responses. Is an essential constituent of several E3 ubiquitin-protein ligase complexes. May have E3 ubiquitin-protein ligase activity and promote \'Lys-63\'-linked ubiquitination of target proteins. Inhibits activation of NF-kappa-B in response to LTBR stimulation. Inhibits TRAF2-mediated activation of NF-kappa-B. Down-regulates proteolytic processing of NFKB2, and thereby inhibits non-canonical activation of NF-kappa-B. Promotes ubiquitination and proteasomal degradation of MAP3K14.

Aliases: CAP 1 antibody, CAP-1 antibody, CAP1 antibody, CD40 associated protein 1 antibody, CD40 binding protein antibody, CD40 bp antibody, CD40 receptor associated factor 1 antibody, CD40 receptor-associated factor 1 antibody, CD40-binding protein antibody, CD40BP antibody, CRAF 1 antibody, CRAF1 antibody, IIAE5 antibody, LAP 1 antibody, LAP1 antibody, LMP1 associated protein antibody, LMP1-associated protein 1 antibody, MIPT3 antibody, TNF receptor associated factor 3 antibody, TRAF 3 antibody, Traf3 antibody, TRAF3\_HUMAN antibody

UniProt:

Q13114

Pathways:

NF-kappaB Signaling, Apoptosis, TLR Signaling, Activation of Innate immune Response, Hepatitis C, Toll-Like Receptors Cascades

#### **Application Details**

Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.