antibodies -online.com







anti-TANK antibody (AA 4-98) (FITC)



Go to Product page

()	11/	IN	/ie	A .
	/ // 	۱ ات	/ (−	' \/\/

Quantity:	100 μg
Target:	TANK
Binding Specificity:	AA 4-98
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TANK antibody is conjugated to FITC
Application:	Please inquire

Product Details

Immunogen:	Recombinant Human TRAF family member-associated NF-kappa-B activator protein (4-98AA)	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	>95%, Protein G purified	

Target Details

Target:	TANK	
Alternative Name:	TANK (TANK Products)	
Background:	Background: Adapter protein involved in I-kappa-B-kinase (IKK) regulation which constituti	
	binds TBK1 and IKBKE playing a role in antiviral innate immunity. Acts as a regulator of TRAF	

function by maintaining them in a latent state. Blocks TRAF2 binding to LMP1 and inhibits LMP1-mediated NF-kappa-B activation. May control negatively TRAF2-mediated NF-kappa-B activation signaled by CD40, TNFR1 and TNFR2.

Aliases: I TRAF antibody, I-TRAF antibody, ITRAF antibody, Tank antibody, TANK_HUMAN antibody, TRAF family member associated NF KAPPA B activator antibody, TRAF family member associated NFKB activator antibody, TRAF family member-associated NF-kappa-B activator antibody, TRAF interacting protein antibody, TRAF interacting protein TANK isoform a antibody, TRAF interacting protein TANK isoform b antibody, TRAF-interacting protein antibody, TRAF2 antibody

UniProt: Q92844

Pathways: p53 Signaling, TLR Signaling, Activation of Innate immune Response

Application Details

Restrictions: For Research Use only

Handling

Format:	Liquid	
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C,-80 °C	
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.	