antibodies

Datasheet for ABIN7172365 anti-TRAPPC6A antibody (AA 1-173)

Image



Overview

Quantity:	100 µg
Target:	TRAPPC6A
Binding Specificity:	AA 1-173
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TRAPPC6A antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human Trafficking protein particle complex subunit 6A protein (1-173AA)
Isotype:	lgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details


Target:	TRAPPC6A
Alternative Name:	TRAPPC6A (TRAPPC6A Products)
Background:	Background: May play a role in vesicular transport during the biogenesis of melanosomes.
	Aliases: TRAPPC6A antibody, HSPC289 antibody, Trafficking protein particle complex subunit

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7172365 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

Target Details

Target Details	
	6A antibody, TRAPP complex subunit 6A antibody
UniProt:	075865
Application Details	
Application Notes:	Recommended dilution: IHC:1:200-1:500,
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Format: Buffer:	Liquid Preservative: 0.03 % Proclin 300
	Preservative: 0.03 % Proclin 300
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Buffer: Preservative:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4 ProClin
Buffer: Preservative:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4 ProClin This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be

Images



Immunohistochemistry

Image 1. IHC image of ABIN7172365 diluted at 1:400 and staining in paraffin-embedded human breast cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.