antibodies - online.com











-						
O	V	e	rv	1	е	W

Quantity:	100 μL
Target:	VGLL1
Binding Specificity:	AA 139-250
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This VGLL1 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human Transcription cofactor vestigial-like protein 1 protein (139-250AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	VGLL1
Alternative Name:	VGLL1 (VGLL1 Products)
Background:	Background: May act as a specific coactivator for the mammalian TEFs.
	Aliases: Protein TONDU antibody, TDU antibody, TONDU antibody, Transcription cofactor

Target Details

vestigial like protein 1 antibody, Transcription cofactor vestigial-like protein 1 antibody, Vestigial
like 1 antibody, vestigial like 1 homolog (Drosophila) antibody, Vgl-1 antibody, VGL1 antibody,
VgII1 antibody, VGLL1_HUMAN antibody

UniProt: Q99990


Application Details

Application Notes:	Recommended dilution: IHC:1:200-1:500,
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Immunohistochemistry

Image 1. IHC image of ABIN7172411 diluted at 1:300 and staining in paraffin-embedded human placenta tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunohistochemistry

Image 2. IHC image of ABIN7172411 diluted at 1:300 and staining in paraffin-embedded human glioma performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.