antibodies -online.com





anti-TEFM antibody (AA 36-165)





Go to Product page

\sim			
	IV/E	۱//۱۲	$I \cap V$

Quantity:	100 μg
Target:	TEFM
Binding Specificity:	AA 36-165
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TEFM antibody is un-conjugated
Application:	Immunohistochemistry (IHC), ELISA

Product Details

Immunogen:	Recombinant Human Transcription elongation factor, mitochondrial protein (36-165AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	TEFM	
Alternative Name:	TEFM (TEFM Products)	
Background:	Background: Transcription elongation factor which increases mitochondrial RNA polymerase processivity. Regulates transcription of the mitochondrial genome, including genes important	

Target Details

UniProt:

for the oxidative phosphorylation machinery.
Aliases: TEFM antibody, C17orf42 antibody, Transcription elongation factor antibody,
mitochondrial antibody
0960E5

Application Details

Application Notes:	Recommended dilution: IHC:1:200-1:500,
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be


Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Immunohistochemistry

Image 1. IHC image of ABIN7172441 diluted at 1:400 and staining in paraffin-embedded human lymph node tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.