
# antibodies .- online.com







### Overview

Quantity:	100 μg
Target:	TAF8
Binding Specificity:	AA 122-311
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TAF8 antibody is conjugated to FITC
Application:	Please inquire

## **Product Details**

Immunogen:	Recombinant Human Transcription initiation factor TFIID subunit 8 protein (122-311AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

# Target Details

Target:	TAF8
Alternative Name:	TAF8 (TAF8 Products)
Background:	Background: Transcription factor TFIID is one of the general factors required for accurate and regulated initiation by RNA polymerase II. Mediates both basal and activator-dependent

transcription. Plays a role in the differentiation of preadipocyte fibroblasts to adipocytes, however, does not seem to play a role in differentiation of myoblasts. Required for the integration of TAF10 in the TAF complex. May be important for survival of cells of the inner cell mass which constitute the pluripotent cell population of the early embryo (By similarity). Aliases: FLJ32821 antibody, hTAFII43 antibody, Protein taube nuss antibody, TAF(II)43 antibody, taf8 antibody, TAF8 RNA polymerase II, TATA box binding protein (TBP) associated factor, 43 kDa antibody, TAF8\_HUMAN antibody, TAFII-43 antibody, TAFII43 antibody, TATA box binding protein (TBP) associated factor, RNA polymerase II A 45/50 kDa antibody, Taube nuss homolog (mouse) antibody, TBN antibody, TBP associated factor 8 antibody, TBP associated factor TAFII43 antibody, TBP-associated factor, RNA polymerase II, 43 kD antibody, TBP-associated factor 8 antibody, Transcription initiation factor TFIID 43 kDa subunit antibody, Transcription initiation factor TFIID subunit 8 antibody

UniProt: Q7Z7C8

Pathways: Maintenance of Protein Location

## **Application Details**

Restrictions: For Research Use only

### Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.