antibodies -online.com





anti-TGFB1I1 antibody (AA 138-200) (HRP)



\sim	
()\/△	rview
\cup	1 410 44

Overview	
Quantity:	100 μg
Target:	TGFB1I1
Binding Specificity:	AA 138-200
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TGFB1I1 antibody is conjugated to HRP
Application:	ELISA
Product Details	
Immunogen:	Recombinant Human Transforming growth factor beta-1-induced transcript 1 protein (138-
	200AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified
Target Details	
Target:	TGFB1I1
Alternative Name:	TGFB1I1 (TGFB1I1 Products)
Background:	Background: Functions as a molecular adapter coordinating multiple protein-protein

interactions at the focal adhesion complex and in the nucleus. Links various intracellular signaling modules to plasma membrane receptors and regulates the Wnt and TGFB signaling pathways. May also regulate SLC6A3 and SLC6A4 targeting to the plasma membrane hence regulating their activity. In the nucleus, functions as a nuclear receptor coactivator regulating glucocorticoid, androgen, mineralocorticoid and progesterone receptor transcriptional activity. May play a role in the processes of cell growth, proliferation, migration, differentiation and senescence. May have a zinc-dependent DNA-binding activity.

Aliases: Androgen receptor coactivator 55 kDa protein antibody, Androgen receptor coactivator ARA55 antibody, Androgen receptor-associated protein of 55 kDa antibody, ARA55 antibody, Hic-5 antibody, Hydrogen peroxide-inducible clone 5 protein antibody, Tgfb1i1 antibody, TGFI1_HUMAN antibody, Transforming growth factor beta 1 induced transcript 1 antibody, Transforming growth factor beta-1-induced transcript 1 protein antibody, TSC 5 antibody

UniProt: 043294

Pathways: Intracellular Steroid Hormone Receptor Signaling Pathway, VEGF Signaling

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.