
antibodies - online.com







anti-TGFB2 antibody (AA 304-413)



Image



Overview

Quantity:	100 μg
Target:	TGFB2
Binding Specificity:	AA 304-413
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TGFB2 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human Transforming growth factor beta-2 proprotein protein (304-413AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	TGFB2
Alternative Name:	TGFB2 (TGFB2 Products)
Background:	Background: TGF-beta 2 has suppressive effects on interleukin-2 dependent T-cell growth.
	Aliases: BSC-1 cell growth inhibitor antibody, BSC1 cell growth inhibitor antibody, Cetermin

Target Details

antibody, G-TSF antibody, Glioblastoma-derived T-cell suppressor factor antibody, GTSF antibody, LAP antibody, Latency-associated peptide antibody, MGC116892 antibody, MGF antibody, Milk growth factor antibody, Polyergin antibody, TGF-beta-2 antibody, TGFB2 antibody, TGFB2_HUMAN antibody, Transforming growth factor beta 2 antibody

UniProt: P61812

Pathways: Cell-Cell Junction Organization, Production of Molecular Mediator of Immune Response,

Protein targeting to Nucleus

Application Details

Application Notes:	Recommended dilution: IHC:1:200-1:500,


Restrictions: For Research Use only

Handling

Storage Comment:

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C

Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Immunohistochemistry

Image 1. IHC image of ABIN7172822 diluted at 1:400 and staining in paraffin-embedded human breast cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.