antibodies - online.com







anti-TMEM64 antibody (AA 40-118)

Images



Overview

| Quantity: | 100 μg |
|----------------------|--|
| Target: | TMEM64 |
| Binding Specificity: | AA 40-118 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This TMEM64 antibody is un-conjugated |
| Application: | ELISA, Immunohistochemistry (IHC), Western Blotting (WB) |

Product Details

| Immunogen: | Recombinant Human Transmembrane protein 64 protein (40-118AA) |
|-------------------|---|
| Isotype: | IgG |
| Cross-Reactivity: | Human, Mouse |
| Purification: | >95%, Protein G purified |

Target Details

| Target: | TMEM64 |
|-------------------|--|
| Alternative Name: | TMEM64 (TMEM64 Products) |
| Background: | Background: Positively regulates TNFSF11-induced osteoclast differentiation. Acts as a regulator of TNFSF11-mediated Ca(2+) signaling pathways via its interaction with SERCA2 |

Target Details

which is critical for the TNFSF11-induced CREB1 activation and mitochondrial ROS generation necessary for proper osteoclast generation. Association between TMEM64 and SERCA2 in the ER leads to cytosolic Ca (2+) spiking for activation of NFATC1 and production of mitochondrial ROS, thereby triggering Ca (2+) signaling cascades that promote osteoclast differentiation and activation. Negatively regulates osteoblast differentiation and positively regulates adipocyte differentiation via modulation of the canonical Wnt signaling pathway. Mediates the switch in lineage commitment to osteogenesis rather than to adipogenesis in mesenchymal stem cells by negatively regulating the expression, activity and nuclear localization of CTNNB1.

Aliases: TMEM64 antibody, TMM64_HUMAN antibody, Transmembrane protein 64 antibody

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be

UniProt:

Q6YI46

Application Details


Precaution of Use:

| Application Notes: | Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200, |
|--------------------|---|
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Buffer: | Preservative: 0.03 % Proclin 300 |
| | Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4 |

Storage: -20 °C,-80 °C


Storage Comment: Upon receipt, store at -20 °C or -80 °C. Avoid repeated freeze.

handled by trained staff only.



Immunohistochemistry

Image 1. Immunohistochemistry of paraffin-embedded human lung tissue using ABIN7173257 at dilution of 1:100



Western Blotting

Image 2. Western blot All lanes: TMEM64 antibody at 8 μ g/mL + Mouse brain tissue Secondary Goat polyclonal to rabbit IgG at 1/10000 dilution Predicted band size: 40, 13, 15, 34 kDa Observed band size: 40 kDa