# antibodies .- online.com







## anti-TGS1 antibody (AA 713-853) (Biotin)



#### Overview

Quantity:	100 μg
Target:	TGS1
Binding Specificity:	AA 713-853
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TGS1 antibody is conjugated to Biotin
Application:	ELISA

#### **Product Details**

Immunogen:	Recombinant Human Trimethylguanosine synthase protein (713-853AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

### Target Details

Target:	TGS1
Alternative Name:	TGS1 (TGS1 Products)
Background:	Background: Catalyzes the 2 serial methylation steps for the conversion of the 7-
	monomethylguanosine (m7G) caps of snRNAs and snoRNAs to a 2,2,7-trimethylguanosine

(m(2,2,7)G) cap structure. The enzyme is specific for guanine, and N7 methylation must precede N2 methylation. Hypermethylation of the m7G cap of U snRNAs leads to their concentration in nuclear foci, their colocalization with coilin and the formation of canonical Cajal bodies (CBs). Plays a role in transcriptional regulation.

Aliases: Cap specific guanine N2 methyltransferase antibody, Cap-specific guanine-N2 methyltransferase antibody, CLL associated antigen KW 2 antibody, CLL-associated antigen KW-2 antibody, DKFZp762A163 antibody, FLJ22995 antibody, HCA137 antibody, Hepatocellular carcinoma associated antigen 137 antibody, Hepatocellular carcinoma-associated antigen 137 antibody, NCOA6IP antibody, Nuclear receptor coactivator 6 interacting protein antibody, Nuclear receptor coactivator 6-interacting protein antibody, PIMT antibody, PIPMT antibody, PRIP interacting protein PIPMT antibody, PRIP interacting protein with methyltransferase domain antibody, PRIP interacting protein with methyltransferase motif antibody, PRIP-interacting protein with methyltransferase motif antibody, TGS 1 antibody, TgS1 antibody, TGS1\_HUMAN antibody, Trimethylguanosine synthase antibody, Trimethylguanosine synthase homolog (S. cerevisiae) antibody, Trimethylguanosine synthase homolog antibody

UniProt: Q96RS0

Pathways: Mitotic G1-G1/S Phases, Regulation of Lipid Metabolism by PPARalpha, Ribonucleoprotein

Complex Subunit Organization

#### **Application Details**

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

#### Handling

Format:	Liquid
- Cirriati	,40.0
Buffer:	Preservative: 0.03 % Proclin 300
	Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be
	handled by trained staff only.
Storage:	-20 °C,-80 °C

Storage Comment:

Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.