

Datasheet for ABIN7173429
anti-TRIM26 antibody (AA 151-270)[Go to Product page](#)

1 Image

Overview

Quantity:	100 µg
Target:	TRIM26
Binding Specificity:	AA 151-270
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TRIM26 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human Tripartite motif-containing protein 26 protein (151-270AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	TRIM26
Alternative Name:	TRIM26 (TRIM26 Products)
Background:	Background: cytosol, DNA binding, metal ion binding, innate immune response, interferon-gamma-mediated signaling pathway, negative regulation of viral entry into host cell, negative

Target Details

regulation of viral release from host cell, positive regulation of sequence-specific DNA binding
transcription factor activity

Aliases: Acid finger protein antibody, AFP antibody, CAM25547 antibody, RING finger protein 95 antibody, RNF95 antibody, TRI26_HUMAN antibody, TRIM26 antibody, tripartite motif containing 26 antibody, Tripartite motif-containing protein 26 antibody, widely expressed acid zinc finger protein antibody, Zinc finger protein 173 antibody, Zinc finger protein 173, formerly antibody, ZNF173 antibody, ZNF173, formerly antibody

UniProt: [Q12899](#)

Application Details

Application Notes: Recommended dilution: IHC:1:200-1:500,

Restrictions: For Research Use only

Handling

Format: Liquid


Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C, -80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Immunohistochemistry

Image 1. IHC image of ABIN7173429 diluted at 1:300 and staining in paraffin-embedded human endometrial cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.