

Datasheet for ABIN7173439
anti-TRIM35 antibody (AA 1-206)[Go to Product page](#)

3 Images

Overview

Quantity:	100 µg
Target:	TRIM35
Binding Specificity:	AA 1-206
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TRIM35 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF)

Product Details

Immunogen:	Recombinant Human Tripartite motif-containing protein 35 protein (1-206AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	TRIM35
Alternative Name:	TRIM35 (TRIM35 Products)
Background:	Background: May play a role as a tumor suppressor. Implicated in the cell death mechanism. Aliases: 0710005M05Rik antibody, A430106H13Rik antibody, AW046487 antibody,

Target Details

Hemopoietic lineage switch protein 5 antibody, HLS5 antibody, KIAA1098 antibody, Macrophage derived apoptosis inducing RBCC protein antibody, MAIR antibody, MGC17233 antibody, mKIAA1098 antibody, NC8 antibody, Protein MAIR antibody, Protein Nc8 antibody, TRI35_HUMAN antibody, TRIM35 antibody, Tripartite motif containing 35 antibody, Tripartite motif containing protein 35 antibody, Tripartite motif-containing protein 35 antibody

UniProt: [Q9UPQ4](#)

Application Details

Application Notes: Recommended dilution: IHC:1:200-1:500, IF:1:50-1:200,

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4


Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C


Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images




Immunofluorescence

Image 1. Immunofluorescence staining of Hela cells with ABIN7173439 at 1:133, counter-stained with DAPI. The cells were fixed in 4 % formaldehyde, permeabilized using 0.2 % Triton X-100 and blocked in 10 % normal Goat Serum. The cells were then incubated with the antibody overnight at 4 °C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemistry

Image 2. IHC image of ABIN7173439 diluted at 1:400 and staining in paraffin-embedded human glioma performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunohistochemistry

Image 3. IHC image of ABIN7173439 diluted at 1:400 and staining in paraffin-embedded human small intestine tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.