antibodies -online.com







anti-TNFAIP8 antibody (AA 1-198)





0 -	4.4	Prod	400		
1-()	10	Pron	ויאוו	nac	\Box
\circ	LO.	1 100	uot	Duc	

\sim							
	1//	\Box	r۱	/ [\bigcirc	1	٨,

Quantity:	100 μL	
Target:	TNFAIP8	
Binding Specificity:	AA 1-198	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This TNFAIP8 antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)	

Product Details

Immunogen:	Recombinant Human Tumor necrosis factor alpha-induced protein 8 protein (1-198AA)	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	Antigen Affinity Purified	

Target Details

Target:	TNFAIP8	
Alternative Name:	TNFAIP8 (TNFAIP8 Products)	
Background:	Background: Acts as a negative mediator of apoptosis and may play a role in tumor	
	progression. Suppresses the TNF-mediated apoptosis by inhibiting caspase-8 activity but not	

Target Details

the processing of procaspase-8, subsequently resulting in inhibition of BID cleavage and caspase-3 activation.

Aliases: GG2 1 antibody, Head and neck tumor and metastasis-related protein antibody, MDC-3.13 antibody, NDED antibody, NF-kappa-B-inducible DED-containing protein antibody, SCC S2 antibody, SCC-S2 antibody, SCC-S2 antibody, TFIP8_HUMAN antibody, TNF alpha-induced protein 8 antibody, TNF induced protein antibody, TNF-induced protein GG2-1 antibody, TNFAIP 8 antibody, TNFAIP8 antibody, Tumor necrosis factor alpha-induced protein 8 antibody


UniProt:

095379

Application Details


Application Notes:	Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200,		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		

Format:	Liquid
Buffer:	PBS with 0.02 % sodium azide, 50 % glycerol, pH 7.3.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.




Immunohistochemistry

Image 1. Immunohistochemistry of paraffin-embedded human breast cancer using ABIN7173776 at dilution of 1:100



Western Blotting

Image 2. Western blot All lanes: TNFAIP8 antibody at $2.32~\mu$ g/mL Lane 1: A431 whole cell lysate Lane 2: PC-3 whole cell lysate Secondary Goat polyclonal to rabbit IgG at 1/10000 dilution Predicted band size: 24, 23, 22, 25 kDa Observed band size: 22 kDa



Immunohistochemistry

Image 3. Immunohistochemistry of paraffin-embedded human colon cancer using ABIN7173776 at dilution of 1:100