



[Go to Product page](#)

Datasheet for ABIN7173778

## anti-TNFAIP8L1 antibody (AA 1-186)

### 1 Image

#### Overview

Quantity:	100 µL
Target:	TNFAIP8L1
Binding Specificity:	AA 1-186
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TNFAIP8L1 antibody is un-conjugated
Application:	ELISA, Immunofluorescence (IF)

#### Product Details

Immunogen:	Recombinant Human Tumor necrosis factor alpha-induced protein 8-like protein 1 protein (1-186AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

#### Target Details

Target:	TNFAIP8L1
Alternative Name:	TNFAIP8L1 ( <a href="#">TNFAIP8L1 Products</a> )
Background:	Background: Acts as a negative regulator of mTOR activity.

## Target Details

Aliases: TNFAIP8L1 antibody, Tumor necrosis factor alpha-induced protein 8-like protein 1 antibody, TIPE1 antibody, TNF alpha-induced protein 8-like protein 1 antibody, TNFAIP8-like protein 1 antibody, Oxidative stress-regulated gene-beta antibody, Oxy-beta antibody

UniProt: [Q8WVP5](#)

## Application Details

Application Notes: Recommended dilution: IF:1:20-1:200,

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4


Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

## Images



### Immunofluorescence

**Image 1.** Immunofluorescence staining of HepG2 cells with ABIN7173778 at 1:50, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).