

Datasheet for ABIN7173832

anti-TNFRSF1B antibody (AA 56-210) (HRP)



Go to Product page

_				
()	ve	r\/		۸ /
	$^{\prime}$ $^{\prime}$: I V	\Box	٧V

Overview		
Quantity:	100 μg	
Target:	TNFRSF1B	
Binding Specificity:	AA 56-210	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This TNFRSF1B antibody is conjugated to HRP	
Application:	ELISA	
Product Details		
Immunogen:	Recombinant Human Tumor necrosis factor receptor superfamily member 10B protein (56-	
	210AA)	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	>95%, Protein G purified	
Target Details		
Target:	TNFRSF1B	
Alternative Name:	TNFRSF1B (TNFRSF1B Products)	
Background:	Background: Receptor for the cytotoxic ligand TNFSF10/TRAIL. The adapter molecule FADD	

recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis. Promotes the activation of NF-kappa-B. Essential for ER stress-induced apoptosis.

Aliases: rotein antibody, Apoptosis inducing protein TRICK2A/2B antibody, Apoptosis inducing receptor TRAIL R2 antibody, CD262 antibody, CD262 antigen antibody, Cytotoxic TRAIL receptor 2 antibody, Death domain containing receptor for TRAIL/Apo 2L antibody, Death receptor 5 antibody, DR5 antibody, KILLER antibody, KILLER/DR5 antibody, OTTHUMP00000123492 antibody, OTTHUMP00000123493 antibody, p53 regulated DNA damage inducible cell death receptor(killer) antibody, TNF related apoptosis inducing ligand receptor 2 antibody, TNF-related apoptosis-inducing ligand receptor 2 antibody, TNFRSF10B antibody, TR10B_HUMAN antibody, TRAIL R2 antibody, TRAIL receptor 2 antibody, TRAIL-R2 antibody, TRICK2 antibody, TRICK2 antibody, TRICK2B antibody, TRICK2B antibody, TRICKB antibody, Tumor necrosis factor receptor like protein ZTNFR9 antibody, Tumor necrosis factor receptor superfamily, member 10B antibody, ZTNFR9 antibody

UniProt: 014763

Pathways: NF-kappaB Signaling, Apoptosis, Cellular Response to Molecule of Bacterial Origin, Hepatitis C,

Ubiquitin Proteasome Pathway

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Format:	Liquid	
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C,-80 °C	

Storage Comment:

Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.