

Datasheet for ABIN7173864
anti-TNFRSF19 antibody (AA 276-415)[Go to Product page](#)

2 Images

Overview

Quantity:	100 µg
Target:	TNFRSF19
Binding Specificity:	AA 276-415
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TNFRSF19 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF)

Product Details

Immunogen:	Recombinant Human Tumor necrosis factor receptor superfamily member 19 protein (276-415AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	TNFRSF19
Alternative Name:	TNFRSF19 (TNFRSF19 Products)
Background:	Background: Can mediate activation of JNK and NF-kappa-B. May promote caspase-

Target Details

independent cell death.

Aliases: AL023044 antibody, AW123854 antibody, TAJ alpha antibody, TAJ antibody, TAJ-alpha antibody, TNFRSF19 antibody, TNFRSF19 tumor necrosis factor receptor superfamily, member 19 antibody, TNFR19_HUMAN antibody, Toxicity and JNK inducer antibody, TRADE antibody, TROY antibody, Tumor necrosis factor receptor superfamily member 19 antibody, UNQ1888/PRO4333 antibody

UniProt: [Q9NS68](#)

Application Details

Application Notes: Recommended dilution: WB:1:500-1:5000, IF:1:50-1:200,

Restrictions: For Research Use only

Handling

Format: Liquid


Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: -20 °C, -80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Immunofluorescence

Image 1. Immunofluorescence staining of HepG2 cells with ABIN7173864 at 1:200, counter-stained with DAPI. The cells were fixed in 4 % formaldehyde, permeabilized using 0.2 % Triton X-100 and blocked in 10 % normal Goat Serum. The cells were then incubated with the antibody overnight at 4 °C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Western Blotting

Image 2. Western Blot Positive WB detected in: PC-3 whole cell lysate All lanes: TNFRSF19 antibody at 2.36 µg/mL Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 47, 46, 31 kDa Observed band size: 47 kDa