antibodies -online.com





anti-LTBR antibody (AA 31-224)

3 Images



Go to Product page

· ·	
$\bigcap \bigcup \bigcap \bigcup \bigcup$	i

Overview	
Quantity:	100 μg
Target:	LTBR
Binding Specificity:	AA 31-224
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LTBR antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF)
Product Details	
Immunogen:	Recombinant Human Tumor necrosis factor receptor superfamily member 3 protein (31-
	224AA)
Isotype:	lgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified
Target Details	
Target:	LTBR
Alternative Name:	LTBR (LTBR Products)
Background:	CD18 antibody, D12S370 antibody, LT beta R antibody, LTBETAR antibody, Ltbr antibody,

Lymphotoxin B receptor antibody, Lymphotoxin beta receptor (TNFR superfamily, member 3) antibody, Lymphotoxin beta receptor antibody, Lymphotoxin-beta receptor antibody, TNF R III antibody, TNF-RIII antibody, TNFCR antibody, TNFR RP antibody, TNFR superfamily member 3 antibody, TNFR-III antibody, TNFR2 RP antibody, TNFR2RP antibody, TNFR3 antibody, TNFRII antibody, TNFRRP antibody, TNFRSF 3 antibody, TNFRSF3 antibody, TNR3_HUMAN antibody, Tumor necrosis factor C receptor antibody, Tumor necrosis factor receptor 2 related protein antibody, Tumor necrosis factor receptor 2-related protein antibody, Tumor necrosis factor receptor superfamily member 3 precursor antibody, Tumor necrosis factor receptor type III antibody

UniProt:

P36941

Pathways:

NF-kappaB Signaling


Application Details


Application Notes:	Recommended dilution: IHC:1:20-1:200, IF:1:50-1:200,
Application Notes.	1.200, ii .1.30 1.200,


Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.







Immunofluorescence

Image 1. Immunofluorescence staining of Hela cells with ABIN7173883 at 1:60, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).

Immunohistochemistry

Image 2. IHC image of ABIN7173883 diluted at 1:180 and staining in paraffin-embedded human breast cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

Immunohistochemistry

Image 3. IHC image of ABIN7173883 diluted at 1:180 and staining in paraffin-embedded human lung cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.