

Datasheet for ABIN7174075

Overview

Target:

Alternative Name:

Background:

anti-ROR2 antibody (AA 752-926) (HRP)

ROR2

ROR2 (ROR2 Products)



Quantity:	100 μg
Target:	ROR2
Binding Specificity:	AA 752-926
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ROR2 antibody is conjugated to HRP
Application:	ELISA
Product Details	
Immunogen:	Recombinant Human Tyrosine-protein kinase transmembrane receptor ROR2 protein (752-
	926AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified
Target Details	

Background: Tyrosine-protein kinase receptor which may be involved in the early formation of

the chondrocytes. It seems to be required for cartilage and growth plate development (By similarity). Phosphorylates YWHAB, leading to induction of osteogenesis and bone formation (PubMed:17717073). In contrast, has also been shown to have very little tyrosine kinase activity in vitro. May act as a receptor for wnt ligand WNT5A which may result in the inhibition of WNT3A-mediated signaling (PubMed:25029443).

Aliases: BDB antibody, BDB1 antibody, Brachydactyly type B antibody, EC 2.7.10.1 antibody, MGC163394 antibody, Neurotrophic tyrosine kinase antibody, Neurotrophic tyrosine kinase, receptor related 2 antibody, NTRKR2 antibody, Receptor tyrosine kinase-like orphan receptor 2 antibody, receptor-related 2 antibody, ROR2 antibody, ROR2_HUMAN antibody, Tyrosine protein kinase transmembrane receptor ROR2 antibody, Tyrosine-protein kinase transmembrane receptor ROR2 antibody

UniProt: Q01974

Pathways: RTK Signaling, WNT Signaling

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Format:

Liquid

Buffer:

Preservative: 0.03 % Proclin 300

Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative:

ProClin

Precaution of Use:

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage:

-20 °C,-80 °C

Upon receipt, store at -20 °C or -80 °C. Avoid repeated freeze.