

Datasheet for ABIN7174119 anti-SHP1 antibody (AA 1-597) (HRP)



Go to Froduct page

	er		

Quantity:	100 μg		
Target:	SHP1 (PTPN6)		
Binding Specificity:	AA 1-597		
Reactivity:	Human		
Host:	Rabbit		
Clonality:	Polyclonal		
Conjugate:	This SHP1 antibody is conjugated to HRP		
Application:	ELISA		
Product Details			
Immunogen:	Recombinant Human Tyrosine-protein phosphatase non-receptor type 6 protein (1-597AA)		
Isotype:	IgG		
Cross-Reactivity:	Human		
Purification:	>95%, Protein G purified		
Target Details			
Target:	SHP1 (PTPN6)		
Alternative Name:	PTPN6 (PTPN6 Products)		
Background:	Background: Modulates signaling by tyrosine phosphorylated cell surface receptors such as KIT and the EGF receptor/EGFR. The SH2 regions may interact with other cellular components to		

modulate its own phosphatase activity against interacting substrates. Together with MTUS1, induces UBE2V2 expression upon angiotensin II stimulation. Plays a key role in hematopoiesis. Aliases: 70Z-SHP antibody, EC 3.1.3.48 antibody, HCP antibody, HCPH antibody, Hematopoietic cell phosphatase antibody, Hematopoietic cell protein tyrosine phosphatase antibody, Hematopoietic cell protein-tyrosine phosphatase antibody, HPTP1C antibody, Protein tyrosine phosphatase 1C antibody, Protein tyrosine phosphatase 1C antibody, Protein-tyrosine phosphatase 1C antibody, protein-tyrosine phosphatase SHP1 antibody, Protein-tyrosine phosphatase SHP-1 antibody, PTP1C antibody, PTP1C antibody, PTP1C antibody, PTP1C antibody, PTP1C antibody, SHP1L antibody, SHP1L antibody, SHP1L antibody, SHP1L antibody, SHP1L antibody, Tyrosine-protein phosphatase non-receptor type 6 antibody

UniProt:

P29350

Pathways:

JAK-STAT Signaling, TCR Signaling, TLR Signaling, Nuclear Receptor Transcription Pathway,
Positive Regulation of Peptide Hormone Secretion, Steroid Hormone Mediated Signaling
Pathway, Response to Growth Hormone Stimulus, Regulation of Leukocyte Mediated Immunity,
CXCR4-mediated Signaling Events, Signaling Events mediated by VEGFR1 and VEGFR2, BCR
Signaling

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.		
Storage:	-20 °C,-80 °C		
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.		