



[Go to Product page](#)

Datasheet for ABIN7174229  
**anti-Ubiquilin 2 antibody (AA 473-542) (Biotin)**

### Overview

Quantity:	100 µg
Target:	Ubiquilin 2 (UBQLN2)
Binding Specificity:	AA 473-542
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Ubiquilin 2 antibody is conjugated to Biotin
Application:	ELISA

### Product Details

Immunogen:	Recombinant Human Ubiquilin-2 protein (473-542AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

### Target Details

Target:	Ubiquilin 2 (UBQLN2)
Alternative Name:	UBQLN2 ( <a href="#">UBQLN2 Products</a> )
Background:	Background: Plays an important role in the regulation of different protein degradation mechanisms and pathways including ubiquitin-proteasome system (UPS), autophagy and the

## Target Details

---

endoplasmic reticulum-associated protein degradation (ERAD) pathway. Mediates the proteasomal targeting of misfolded or accumulated proteins for degradation by binding (via UBA domain) to their polyubiquitin chains and by interacting (via ubiquitin-like domain) with the subunits of the proteasome (PubMed:10983987). Plays a role in the ERAD pathway via its interaction with ER-localized proteins FAF2/UBXD8 and HERPUD1 and may form a link between the polyubiquitinated ERAD substrates and the proteasome (PubMed:24215460, PubMed:18307982). Involved in the regulation of macroautophagy and autophagosome formation, required for maturation of autophagy-related protein LC3 from the cytosolic form LC3-I to the membrane-bound form LC3-II and may assist in the maturation of autophagosomes to autolysosomes by mediating autophagosome-lysosome fusion (PubMed:19148225, PubMed:20529957). Negatively regulates the endocytosis of GPCR receptors: AVPR2 and ADRB2, by specifically reducing the rate at which receptor-arrestin complexes concentrate in clathrin-coated pits (CCPs) (PubMed:18199683).

Aliases: UBQLN2 antibody, N4BP4 antibody, PLIC2 antibody, HRIHFB2157Ubiquilin-2 antibody, Chap1 antibody, DSK2 homolog antibody, Protein linking IAP with cytoskeleton 2 antibody, PLIC-2 antibody, hPLIC-2 antibody, Ubiquitin-like product Chap1/Dsk2 antibody

---

UniProt: [Q9UHD9](#)

## Application Details

---

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.