



[Go to Product page](#)

Datasheet for ABIN7174271

## anti-USP17L5 antibody (C-Term)

### 3 Images

#### Overview

Quantity:	100 µL
Target:	USP17L5
Binding Specificity:	AA 19-235, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This USP17L5 antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (IHC)

#### Product Details

Immunogen:	Recombinant Human Ubiquitin carboxyl-terminal hydrolase 17-like protein 5 protein (19-235AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

#### Target Details

Target:	USP17L5
Alternative Name:	USP17L5 ( <a href="#">USP17L5 Products</a> )
Background:	Background: Deubiquitinating enzyme that removes conjugated ubiquitin from specific proteins to regulate different cellular processes that may include cell proliferation, progression through

## Target Details

the cell cycle, apoptosis, cell migration, and the cellular response to viral infection.

Aliases: USP17L5 antibody, Ubiquitin carboxyl-terminal hydrolase 17-like protein 5 antibody, EC 3.4.19.12 antibody, Deubiquitinating enzyme 17-like protein 5 antibody, Ubiquitin thioesterase 17-like protein 5 antibody, Ubiquitin-specific-processing protease 17-like protein 5 antibody

UniProt: [A8MUK1](#)

## Application Details

Application Notes: Recommended dilution: WB:1:1000-1:5000, IHC:1:200-1:500,

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4


Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C


Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

## Images




### Immunohistochemistry

**Image 1.** IHC image of ABIN7174271 diluted at 1:400 and staining in paraffin-embedded human liver cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



### Western Blotting

**Image 2.** Western Blot Positive WB detected in: HeLa whole cell lysate All lanes: USP17L5 antibody at 1:2000 Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 60 kDa Observed band size: 60 kDa



### Immunohistochemistry

**Image 3.** IHC image of ABIN7174271 diluted at 1:400 and staining in paraffin-embedded human gastric cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.