



[Go to Product page](#)

Datasheet for ABIN7174418  
**anti-UBAC2 antibody (AA 1-157)**

2 Images

### Overview

Quantity:	100 µg
Target:	UBAC2
Binding Specificity:	AA 1-157
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This UBAC2 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

### Product Details

Immunogen:	Recombinant Human Ubiquitin-associated domain-containing protein 2 protein (1-157AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

### Target Details

Target:	UBAC2
Alternative Name:	UBAC2 ( <a href="#">UBAC2 Products</a> )
Background:	Background: Restricts trafficking of FAF2 from the endoplasmic reticulum to lipid droplets. Aliases: UBAC2 antibody, PHGDHL1 antibody, PSEC0110 antibody, Ubiquitin-associated

## Target Details

domain-containing protein 2 antibody, UBA domain-containing protein 2 antibody,  
Phosphoglycerate dehydrogenase-like protein 1 antibody

UniProt: [Q8NBM4](#)

## Application Details

Application Notes: Recommended dilution: IHC:1:200-1:500,

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4


Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C


Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

## Images



### Immunohistochemistry

**Image 1.** IHC image of ABIN7174418 diluted at 1:300 and staining in paraffin-embedded human testis tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



### Immunohistochemistry

**Image 2.** IHC image of ABIN7174418 diluted at 1:300 and staining in paraffin-embedded human colon cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.