Datasheet for ABIN7174950 anti-MYO1G antibody (AA 209-389) (Biotin)

-online.com antibodies



Overview

| Quantity:            | 100 µg                                      |
|----------------------|---|
| Target:              | MY01G                                       |
| Binding Specificity: | AA 209-389                                  |
| Reactivity:          | Human                                       |
| Host:                | Rabbit                                      |
| Clonality:           | Polyclonal                                  |
| Conjugate:           | This MYO1G antibody is conjugated to Biotin |
| Application:         | ELISA                                       |

#### Product Details

| Immunogen:        | Recombinant Human Unconventional myosin-Ig protein (209-389AA) |
|-------------------|--|
| lsotype:          | lgG  |
| Cross-Reactivity: | Human  |
| Purification:     | >95%, Protein G purified                                       |

### Target Details

| Target:           | MY01G   |
|-------------------|---|
| Alternative Name: | MY01G (MY01G Products)  |
| Background:       | Background: Unconventional myosin required during immune response for detection of rare         |
|                   | antigen-presenting cells by regulating T-cell migration. Unconventional myosins are actin-based |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7174950 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

### Target Details

| UniProt: | B0I1T2  |
|----------|---|
|          | MYO1G_HUMAN antibody, myosin IG antibody  |
|          | Aliases: mHag HA-2 antibody, Minor histocompatibility antigen HA-2 antibody, Myo1g antibody,    |
|          | phagocytosis.   |
|          | membrane/cytoskeleton-dependent processes. Involved in Fc-gamma receptor (Fc-gamma-R)           |
|          | pathogen eradication. Also required in B-cells, where it regulates different                    |
|          | thereby enhancing detection of rare antigens during lymph-node surveillance, enabling           |
|          | of T-cell migration by generating membrane tension, enforcing cell-intrinsic meandering search, |
|          | motor molecules with ATPase activity and serve in intracellular movements. Acts as a regulator  |

# Application Details

| Application Notes: | Optimal working dilution should be determined by the investigator. |
|--------------------|--|
| Restrictions:      | For Research Use only  |

# Handling

| Format:            | Liquid  |
|--------------------|---|
| Buffer:            | Preservative: 0.03 % Proclin 300<br>Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4                                |
| Preservative:      | ProClin   |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage:           | -20 °C,-80 °C   |
| Storage Comment:   | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.   |