

[Go to Product page](#)

Datasheet for ABIN7174951
anti-MYO1G antibody (AA 209-389) (FITC)

Overview

| | |
|----------------------|---|
| Quantity: | 100 µg |
| Target: | MYO1G |
| Binding Specificity: | AA 209-389 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This MYO1G antibody is conjugated to FITC |
| Application: | Please inquire |

Product Details

| | |
|-------------------|--|
| Immunogen: | Recombinant Human Unconventional myosin-Ig protein (209-389AA) |
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Purification: | >95%, Protein G purified |

Target Details

| | |
|-------------------|---|
| Target: | MYO1G |
| Alternative Name: | MYO1G (MYO1G Products) |
| Background: | Background: Unconventional myosin required during immune response for detection of rare antigen-presenting cells by regulating T-cell migration. Unconventional myosins are actin-based |

Target Details

motor molecules with ATPase activity and serve in intracellular movements. Acts as a regulator of T-cell migration by generating membrane tension, enforcing cell-intrinsic meandering search, thereby enhancing detection of rare antigens during lymph-node surveillance, enabling pathogen eradication. Also required in B-cells, where it regulates different membrane/cytoskeleton-dependent processes. Involved in Fc-gamma receptor (Fc-gamma-R) phagocytosis.

Aliases: mHag HA-2 antibody, Minor histocompatibility antigen HA-2 antibody, Myo1g antibody, MYO1G_HUMAN antibody, myosin IG antibody

UniProt: [B0I1T2](#)

Application Details

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.