# antibodies - online.com







# anti-Myosin XIX antibody (AA 2-770)



# Image



#### Overview

Quantity:	100 μg
Target:	Myosin XIX (MYO19)
Binding Specificity:	AA 2-770
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Myosin XIX antibody is un-conjugated
Application:	ELISA, Western Blotting (WB)

### **Product Details**

Immunogen:	Recombinant Human Unconventional myosin-XIX protein (2-770AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

# Target Details

Target:	Myosin XIX (MYO19)
Alternative Name:	MYO19 (MYO19 Products)
Background:	Background: Myosins are actin-based motor molecules with ATPase activity. Unconventional
	myosins serve in intracellular movements. Their highly divergent tails are presumed to bind to

#### **Target Details**

membranous compartments, which would be moved relative to actin filaments. MYO19 is involved in mitochondrial motility.

Aliases: FLJ22865 antibody, myo19 antibody, MYO19\_HUMAN antibody, Myohd1 antibody, Myosin head domain containing 1 antibody, Myosin head domain-containing protein 1 antibody, myosin XIX antibody, Myosin-XIX antibody, unconventional myosin-XIX antibody

UniProt: Q96H55


#### **Application Details**

Application Notes:	Recommended dilution: WB:1:200-1:1000,
Restrictions:	For Research Use only

#### Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

#### **Images**



#### **Western Blotting**

Image 1. Western blot All lanes: MYO19 antibody at  $2\,\mu$  g/mL + A549 whole cell lysate Secondary Goat polyclonal to rabbit IgG at 1/10000 dilution Predicted band size: 110, 36, 72, 87 kDa Observed band size: 110 kDa