
antibodies .- online.com





anti-Myosin XVIIIA antibody (AA 1849-2054) (HRP)



Go to Product page

()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL
Target:	Myosin XVIIIA (MYO18A)
Binding Specificity:	AA 1849-2054
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Myosin XVIIIA antibody is conjugated to HRP
Application:	ELISA

Product Details

Immunogen:	Recombinant Human Unconventional myosin-XVIIIa protein (1849-2054AA)	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	>95%, Protein G purified	

Target Details

Target:	Myosin XVIIIA (MYO18A)	
Alternative Name:	MYO18A (MYO18A Products)	
Background:	Background: May link Golgi membranes to the cytoskeleton and participate in the tensile force	
	required for vesicle budding from the Golgi. Thereby, may play a role in Golgi membrane	

trafficking and could indirectly give its flattened shape to the Golgi apparatus (PubMed:19837035, PubMed:23345592). Alternatively, in concert with LURAP1 and CDC42BPA/CDC42BPB, has been involved in modulating lamellar actomyosin retrograde flow that is crucial to cell protrusion and migration (PubMed:18854160). May be involved in the maintenance of the stromal cell architectures required for cell to cell contact (By similarity). Regulates trafficking, expression, and activation of innate immune receptors on macrophages. Plays a role to suppress inflammatory responsiveness of macrophages via a mechanism that modulates CD14 trafficking (PubMed:25965346). Acts as a receptor of surfactant-associated protein A (SFTPA1/SP-A) and plays an important role in internalization and clearance of SFTPA1-opsonized S.aureus by alveolar macrophages (PubMed:16087679, PubMed:21123169). Strongly enhances natural killer cell cytotoxicity (PubMed:27467939). Aliases: MYO18A antibody, CD245 antibody, KIAA0216 antibody, MYSPDZ antibody, Unconventional myosin-XVIIIa antibody, Molecule associated with JAK3 N-terminus antibody, MAJN antibody, Myosin containing a PDZ domain antibody, Surfactant protein receptor SP-R210 antibody, SP-R210 antibody

UniProt:

Q92614

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300
	Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be
	handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.