



[Go to Product page](#)

Datasheet for ABIN7175200  
**anti-VSTM2L antibody (AA 25-204)**

1 Image

### Overview

Quantity:	100 µL
Target:	VSTM2L
Binding Specificity:	AA 25-204
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This VSTM2L antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

### Product Details

Immunogen:	Recombinant Human V-set and transmembrane domain-containing protein 2-like protein (25-204AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Antigen Affinity Purified

### Target Details

Target:	VSTM2L
Alternative Name:	VSTM2L ( <a href="#">VSTM2L Products</a> )
Background:	Background: cytoplasm, extracellular region, negative regulation of neuron apoptotic process

## Target Details

Aliases: C20orf102 antibody, dJ1118M15.2 antibody, Uncharacterized protein C20orf102 precursor antibody, V set and transmembrane domain containing 2 like antibody, V set and transmembrane domain containing protein 2 like protein antibody, V-set and transmembrane domain-containing protein 2-like protein antibody, VSTM 2L antibody, Vstm2l antibody, VTM2L\_HUMAN antibody

UniProt: [Q96N03](#)

## Application Details

Application Notes: Recommended dilution: IHC:1:20-1:200,

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4


Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

## Images



### Immunohistochemistry

**Image 1.** IHC image of ABIN7175200 diluted at 1:100 and staining in paraffin-embedded human colon cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a biotinylated secondary antibody and

visualized using an HRP conjugated SP system.