

Datasheet for ABIN7175387

anti-VEGFR2/CD309 antibody (AA 20-764) (HRP)[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	VEGFR2/CD309 (VEGFR2)
Binding Specificity:	AA 20-764
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This VEGFR2/CD309 antibody is conjugated to HRP
Application:	ELISA

Product Details

Immunogen:	Recombinant Human Vascular endothelial growth factor receptor 2 protein (20-764AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	VEGFR2/CD309 (VEGFR2)
Alternative Name:	KDR (VEGFR2 Products)
Background:	Background: Tyrosine-protein kinase that acts as a cell-surface receptor for VEGFA, VEGFC and VEGFD. Plays an essential role in the regulation of angiogenesis, vascular development,

Target Details

vascular permeability, and embryonic hematopoiesis. Promotes proliferation, survival, migration and differentiation of endothelial cells. Promotes reorganization of the actin cytoskeleton. Isoforms lacking a transmembrane domain, such as isoform 2 and isoform 3, may function as decoy receptors for VEGFA, VEGFC and/or VEGFD. Isoform 2 plays an important role as negative regulator of VEGFA- and VEGFC-mediated lymphangiogenesis by limiting the amount of free VEGFA and/or VEGFC and preventing their binding to FLT4. Modulates FLT1 and FLT4 signaling by forming heterodimers. Binding of vascular growth factors to isoform 1 leads to the activation of several signaling cascades. Activation of PLCG1 leads to the production of the cellular signaling molecules diacylglycerol and inositol 1,4,5-trisphosphate and the activation of protein kinase C. Mediates activation of MAPK1/ERK2, MAPK3/ERK1 and the MAP kinase signaling pathway, as well as of the AKT1 signaling pathway. Mediates phosphorylation of PIK3R1, the regulatory subunit of phosphatidylinositol 3-kinase, reorganization of the actin cytoskeleton and activation of PTK2/FAK1. Required for VEGFA-mediated induction of NOS2 and NOS3, leading to the production of the signaling molecule nitric oxide (NO) by endothelial cells. Phosphorylates PLCG1. Promotes phosphorylation of FYN, NCK1, NOS3, PIK3R1, PTK2/FAK1 and SRC.

Aliases: CD309 antibody, CD309 antigen antibody, EC 2.7.10.1 antibody, Fetal liver kinase 1 antibody, FLK-1 antibody, FLK1 antibody, FLK1, mouse, homolog of antibody, Kdr antibody, Kinase insert domain receptor (a type III receptor tyrosine kinase) antibody, Kinase insert domain receptor antibody, KRD1 antibody, Ly73 antibody, Protein tyrosine kinase receptor FLK1 antibody, Protein-tyrosine kinase receptor flk-1 antibody, soluble VEGFR2 antibody, Tyrosine kinase growth factor receptor antibody, Vascular endothelial growth factor receptor 2 antibody, VEGFR 2 antibody, VEGFR antibody, VEGFR-2 antibody, VEGFR2 antibody, VGFR2_HUMAN antibody

UniProt:	P35968
Pathways:	RTK Signaling , Glycosaminoglycan Metabolic Process , Signaling Events mediated by VEGFR1 and VEGFR2 , Growth Factor Binding , Regulation of long-term Neuronal Synaptic Plasticity , VEGF Signaling

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.