

[Go to Product page](#)

Datasheet for ABIN7175620

**anti-KCNAB1 antibody (AA 287-401) (FITC)**

## Overview

Quantity:	100 µg
Target:	KCNAB1
Binding Specificity:	AA 287-401
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This KCNAB1 antibody is conjugated to FITC
Application:	Please inquire

## Product Details

Immunogen:	Recombinant Human Voltage-gated potassium channel subunit beta-1 protein (287-401AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

## Target Details

Target:	KCNAB1
Alternative Name:	KCNAB1 ( <a href="#">KCNAB1 Products</a> )
Background:	Background: Cytoplasmic potassium channel subunit that modulates the characteristics of the channel-forming alpha-subunits (PubMed:7499366, PubMed:7603988,

## Target Details

PubMed:17156368, PubMed:17540341, PubMed:19713757). Modulates action potentials via its effect on the pore-forming alpha subunits (By similarity). Promotes expression of the pore-forming alpha subunits at the cell membrane, and thereby increases channel activity (By similarity). Mediates closure of delayed rectifier potassium channels by physically obstructing the pore via its N-terminal domain and increases the speed of channel closure for other family members (PubMed:9763623). Promotes the closure of KCNA1, KCNA2 and KCNA5 channels (PubMed:7499366, PubMed:7890032, PubMed:7603988, PubMed:7649300, PubMed:8938711, PubMed:12077175, PubMed:12130714, PubMed:15361858, PubMed:17540341, PubMed:19713757). Accelerates KCNA4 channel closure (PubMed:7890032, PubMed:7649300, PubMed:7890764, PubMed:9763623). Accelerates the closure of heteromeric channels formed by KCNA1 and KCNA4 (PubMed:17156368). Accelerates the closure of heteromeric channels formed by KCNA2, KCNA5 and KCNA6 (By similarity). Isoform KvB1.2 has no effect on KCNA1, KCNA2 or KCNB1 (PubMed:7890032, PubMed:7890764). Enhances KCNB1 and KCNB2 channel activity (By similarity). Binds NADPH, this is required for efficient down-regulation of potassium channel activity (PubMed:17540341). Has NADPH-dependent aldoketoreductase activity (By similarity). Oxidation of the bound NADPH strongly decreases N-type inactivation of potassium channel activity (By similarity).

Aliases: hKvb3 antibody, hKvBeta3 antibody, K(+) channel subunit beta-1 antibody, KCAB1\_HUMAN antibody, KCNA1B antibody, KCNAB1 antibody, Kv-beta-1 antibody, Kvb1.3 antibody, Voltage-gated potassium channel beta-1 subunit antibody, Voltage-gated potassium channel subunit beta-1 antibody

UniProt: [Q14722](#)

## Application Details

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

## Handling

---

Storage: -20 °C, -80 °C

---

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.