antibodies .- online.com







anti-WDR62 antibody (AA 889-1188) (HRP)



| () | 11/0 | K\ / | iew | 1 |
|-----|--------|--------|---------------------|---|
| | \cup | 'I V/I | $\square \vee \vee$ | ı |
| | | | | |

| Quantity: | 100 μg |
|----------------------|--|
| Target: | WDR62 |
| Binding Specificity: | AA 889-1188 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This WDR62 antibody is conjugated to HRP |
| Application: | ELISA |

Product Details

| Immunogen: | Recombinant Human WD repeat-containing protein 62 protein (889-1188AA) | |
|-------------------|--|--|
| Isotype: | IgG | |
| Cross-Reactivity: | Human | |
| Purification: | >95%, Protein G purified | |

Target Details

| Target: | WDR62 | |
|-------------------|--|--|
| Alternative Name: | WDR62 (WDR62 Products) | |
| Background: | Background: Required for cerebral cortical development. Plays a role in neuronal proliferation | |
| | and migration (PubMed:20890278, PubMed:20729831). Plays a role in mother-centriole- | |

Target Details

| Aliases: WDR62 antibody, C19orf14 antibody, WD repeat-containing protein 62 antibody |
|--|
| for centriole duplication (PubMed:26297806). |
| CEP63 through a stepwise assembled complex at the centrosome that recruits CDK2 required |
| dependent centriole duplication, the function seems also to involve CEP152, CDK5RAP2 and |

UniProt:

043379

Application Details

| Application Notes: | Optimal working dilution should be determined by the investigator. | |
|--------------------|--|--|
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Buffer: | Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4 | |

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.