

Datasheet for ABIN7175810
anti-WDR92 antibody (AA 122-357)



[Go to Product page](#)

1 Image

Overview

Quantity:	100 µL
Target:	WDR92
Binding Specificity:	AA 122-357
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This WDR92 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human WD repeat-containing protein 92 protein (122-357AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	WDR92
Alternative Name:	WDR92 (WDR92 Products)
Background:	Background: Seems to act as a modulator of apoptosis. Aliases: FLJ31741 antibody, Monad antibody, OTTHUMP00000205522 antibody, WD repeat

Target Details

domain 92 antibody, WD repeat-containing protein 92 antibody, WD repeat-containing protein
Monad antibody, Wdr92 antibody, WDR92_HUMAN antibody

UniProt: [Q96MX6](#)

Application Details

Application Notes: Recommended dilution: IHC:1:500-1:1000,

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4


Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Immunohistochemistry

Image 1. IHC image of ABIN7175810 diluted at 1:500 and staining in paraffin-embedded human testis tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.