antibodies - online.com







anti-ZNF282 antibody (AA 240-384)



Image



Overview

| Quantity: | 100 μg |
|----------------------|---------------------------------------|
| Target: | ZNF282 |
| Binding Specificity: | AA 240-384 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This ZNF282 antibody is un-conjugated |
| Application: | ELISA, Immunohistochemistry (IHC) |

Product Details

| Immunogen: | Recombinant Human Zinc finger protein 282 protein (240-384AA) |
|-------------------|---|
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Purification: | >95%, Protein G purified |

Target Details

| Target: | ZNF282 |
|-------------------|--|
| Alternative Name: | ZNF282 (ZNF282 Products) |
| Background: | Background: Binds to the U5 repressive element (U5RE) of the human T cell leukemia virus type |
| | I long terminal repeat. It recognizes the 5\'-TCCACCCC-3\' sequence as a core motif and exerts |

Target Details

| a strong repressive effect on HTLV-I LTR-mediated expression. |
|---|
| Aliases: ZNF282 antibody, HUB1Zinc finger protein 282 antibody, HTLV-I U5RE-binding protein |
| 1 antibody, HUB-1 antibody |
| |

UniProt:

Q9UDV7


Application Details

| Application Notes: | Recommended dilution: IHC:1:200-1:500, |
|--------------------|--|
| Restrictions: | For Research Use only |

Handling

| Format: | Liquid |
|--------------------|---|
| Buffer: | Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4 |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | -20 °C,-80 °C |
| Storage Comment: | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |

Images



Immunohistochemistry

Image 1. IHC image of ABIN7176211 diluted at 1:400 and staining in paraffin-embedded human brain tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30 min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.